



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME V

New names, and the final members of new combinations, are in **bold face type**

- Abies, 233, 235; balsamea, 233-239; pectinata, 151, 152
 Abortiporus distortus, 314
 Acer Negundo, 264
 Acrosporium, 57, 58, 61; **compactum**, 58; **Euonymi-japonici**, 58; **Gossypii**, 59, 61; **hyalina**, 58; **leuconium**, 58; monilioides, 57, 58; **obductum**, 58; **pirinum**, 58; **Tuckeri**, 58
 Aecia, Internal, 303
 Agaricaceae of the Pacific Coast, The—IV. New species of Clitocybe and Melanoleuca, 206
 Agaricaceae of tropical North America, The—VI, 18
 Agaricus, 81; agglutinatius, 83; alutaceus, 310; asper, 75; baccatus, 84; badius, 82; bulbosus, 74; bulbosus vernus, 74; cacaophyllus, 27; Caesareus, 73; campestris, 72, 169, 225; camphoratus, 307; Ceciliae, 86; chlorinosmus, 78; chrysopellus, 23; chrysotrichus, 21; circellatus, 306; citrinus, 74; daucipes, 84; deliciosus, 305; echinocephalus, 77; excelsus, 80; flocculentus, 36; Frostianus, 76; fulvus, 82; helvoliceps, 20; hyalinus, 82; inauratus, 86; insulsus, 306; lactifluus dulcis, 307; Listeri, 307; macro-mastes, 36; maculatus, 80; magnificus, 75; martianus, 36; mitissimus, 307; monticulosus, 77; musaecola, 27; muscarius major, 77; muscarius minor, 76; myodes, 75; nigricans, 308; nitidus, 80; nivalis, 84; onustus, 77; pantherinus, 80; peregrinus, 26; phalloides, 74; piperatus, 307; plumbeus, 82; polypyraxis, 77; porphyrius, 81; praetorius, 85; pustulatus, 75; pyrrhus, 28; pyrrhus leiosporus, 28; Ravenelii, 77; recutitus, 81; ricensis, 26; rubens, 75; rubescens, 75; russuloides, 77; scobifer, 35; scrobiculatus, 305; soleatus, 83; solitarius, 77; spissus, 81; spretus, 73; stramineus, 74; strangulatus, 85; strobiliformis, 77, 80; subdulcis, 307; theiogalus, 306; torminosus, 305; trivialis, 306; vaginatus, 82; vellereus, 307; verrucosus, 75; virosus, 81; volvatus, 83; zonarius, 306
 Akebia quinata, 248
 Aleuria fulgens, 302
 Algal host, The lichen and its. The nature and classification of lichens—II, 97
 Alternaria, 279
 Alysidium, 47; fulvum, 47, 48
 Amanita, 72, 226, 229, 230, 232; abrupta, 79; Amici, 77; ampla, 80; aspera, 75; bisporigera, 74; Caesarea, 73, 225, 316; candida, 78; cinereoconia, 78; citrina alba, 80; cothurnata, 74, 96; crenulata, 77; elliptosperma, 79; elongata, 79; farinosa, 84; flavoconia, 76; flavo-rubens, 79; flavorubescens, 76, 95; floccocephala, 74; Frostiana, 96; glabriceps, 79; junquillea, 77, 80; lignophila, 74; livida, 82; magnivelaris, 80; Mappa, 74; Morrisii, 75; multisquamosa, 78; muscaria, 75, 93, 94, 224-227, 229-232, 316; muscaria coccinea, 83; pantherina, 74, 75; **Peckiana**, 67; pellucida, 73; phalloides, 93, 225, 316; prairiicola, 78; pubescens, 85; radi-cata, 78; recutita, 74; rubescens, 75; spadicea, 82; submaculata, 80; umbrina, 81; velatipes, 75; verna, 74, 79, 80; vernalis, 77; virosa, 74
 Amanitas of eastern North America, The, 72
 Amanitopsis, 81, 84; adnata, 84; agglutinata, 83; albocreata, 84; bac-cata, 83; farinosa, 84; hyperborea, 85; parcovolvata, 83; pubescens, 85; pulverulenta, 85; pusilla, 83; strangulata, 85, 86; vaginata, 82; volvata, 83, 94
 American mycological literature, In-dex to, 41, 91, 182, 251, 284, 317
 Amerosporium Vanilla, 40
 Amsinckia intermedia, 313
 Anabena, 131
 Analytic work whether colonies of the chestnut blight fungus originate from pycnosporos or ascospores, A method of determining in, 274
 Anthurus, 268, 272; borealis, 268, 269

- Armillaria mellea*, 38, 314, 316; *mucida*, 38; *subannulata*, 216
Arnoldia, 124, 137
Arthonia dispersa, 139; *punctiformis*, 139; *radiata*, 109, 138; *vulgaris*, 122
Arthopyrenia, 110; *cerasi*, 139; *punctiformis*, 110; *rhyponta*, 139
Arthur, J. C., *Uredinales* on *Carex* in North America, 240
Ascospores, A method of determining in analytic work whether colonies of the chestnut blight fungus originate from pycnospores or, 274
Aseroe, 268
Aspergillus, 46, 47
Aspidium *Thelypteris*, 236, 239
Aster, 242, 264
Auriscalpium, 298
- Bad year for fleshy fungi, A, 315
Banker, H. J., Type studies in the *Hydnaceae*—III. The genus *Sarcodon*, 12; IV. The genus *Phellodon*, 62; V. The genus *Hydnellum*, 194; VI. The genera *Creolophus*, *Echinodontium*, *Gloiodon*, and *Hydnodon*, 293
Barlaea fulgens, 302
Barya, 111
Basidia, 105
Betula odorata, 89
Biatorina Bouteillii, 151; *synthesa*, 150
Bilimbia, 105
Bjerkandera, 313
Blight fungus originate from pycnospores or ascospores, A method of determining in analytic work whether colonies of the chestnut, 274
Boletus, 176; *americanus*, 3; *flavidus*, 4
Botrydina vulgaris, 114
Botrytis, 46
Buellia parasema, 148; *punctiformis*, 122, 123
Burlingham, G. S., The *Lactarieae* of the Pacific Coast, 305
Byssonectria, 177
- Caeoma Abietis-canadensis*, 238; *nitens*, 281, 282
Calicium, 105; *curtum*, 123; *parietinum*, 123; *trachelinum*, 123
Caloscypha, 299
Calospora Vanillae, 39
Caloscypha fulgens, 302
Calvatia cyathiformis, 316; *maxima*, 316
Cantharellus, 261, 262; *brevipes*, 261, 262; *clavatus*, 261, 263; *floccosus*, 262
Cantharellus brevipes and *Cantharellus clavatus*, The identity of, 261
Cantharellus clavatus, The identity of *Cantharellus brevipes* and, 261
Carex, 240–244
Carex in North America, *Uredinales* on, 240
Catillaria denigrata, 123; *prasina*, 123
Cephaleuros Henningsii, 40
Ceriumyces auriporus, 2; *bicolor*, 4; *communis*, 260; *fumosipes*, 259; *illudens*, 260; *Peckii*, 5; *speciosus*, 5; *subglabripes*, 4
Chaenotheca, 113; *chrysocephala*, 113
Chalara mycoderma, 45
Chamonixia, 313
Chanterel cinnabarinus, 258; *minor*, 257
Chestnut blight fungus originate from pycnospores or ascospores, A method of determining in analytic work whether colonies of the, 274
Chionanthus virginica, 248
Chlorella, 129; *viridis*, 131
Chlorococcum, 119, 125, 127, 145, 146; *humicola*, 109, 115, 117, 118, 124, 125, 129–131
Chondromyces aurantiacum, 60
Chromocrea, 179
Chromocreopsis, 179
Chroococcus, 126, 131
Cladonia, 131, 150; *pyxidata*, 137
Cladothrix, 125
Clark, E. D., & Smith, C. S., Toxicological studies on the mushrooms *Clitocybe illudens* and *Inocybe infida*, 224
Classification of lichens, The nature and—II. The lichen and its algal host, 97
Clathrella chrysomycelina, 268; *Clathrus*, 313
Clathrogaster, 313
Clathrus, 267, 268, 271; *cancellatus*, 267
Clavaria mucida, 115; *pistillaris*, 262, 263
Claviceps, 178
Climacodon, 293
Clitocybe, 207, 210, 225, 226, 229–231; *albicastanea*, 206; *albiformis*, 206; *atrialba*, 207, 208; *avellaneialba*, 207; *brunnescens*, 208; *cuticolor*, 208; *cyathiformis*, 208; *dealbata sudorifica*, 225; *griseifolia*, 208; *Harperi*, 209; *hondensis*, 209; *illudens*, 225–227, 229–232; *multiceps*, 225, 230, 232; *murinifolia*, 210; *oculata*, 210; *oreades*, 210; *oregonensis*, 211; *Peckii*, 211; *sinopica*, 212; *stipitata*, 211; *subcandicans*, 212; *subfumosipes*, 212; *subinversa*, 212; *variabilis*, 213; *variabilis*

- brevipes**, 213; **violaceifolia**, 213; **washingtonensis**, 214
Clitocybe and *Melanoleuca*, New species of. The Agaricaceae of the Pacific Coast—IV, 206
Clitocybe illudens and *Inocybe infida*, Toxicological studies on the mushrooms, 224
Clitopilus, 314; *abortivus*, 314
Coast, The Agaricaceae of the Pacific—IV. New species of *Clitocybe* and *Melanoleuca*, 206
Coast, The Lactarieae of the Pacific, 305
Coccoloba uvifera, 246, 247
Cocomyxa subellipsoidea, 114
Collema, 112, 115, 119, 120, 124; *microphyllum*, 123; *pulposum*, 107, 130
Colletotrichum, 171
Collybia, 216, 217; *platyphylla*, 207, 208; **subdecumbens**, 68; **truncata**, 68
Colonies of the chestnut blight fungus originate from pycnospores or ascospores, A method of determining in analytic work whether, 274
Color guide, Ridgway's new, 172
Colus Garciae, 268; *hirudinosus*, 268
Conrad, H. S., The structure of *Sinibulum sphaerocephalum*, 264
Conida, 114; *punctatella*, 114; *rubescens*, 114
Coniothyrium Chionanthi, 248
Conocybe tener, 36
Cookeina, 185-187; *Afzelii*, 189; **Colensoi**, 185, 187, 191, 193; *Hindsii*, 189; *insititia*, 185, 187, 190, 193; *sulcipes*, 187, 189, 190, 192, 193; *Tricholoma*, 186-188, 193
Coprinus, 168; *atramentarius*, 168, 316; *comatus*, 168, 315
Cora, 115, 125, 126
Cordyceps, 178, 179; *Cockerellii*, 179
Coriolus, 313; *prolificans*, 288; *versicolor*, 287
Corticium salmonicolor, 39
Craterellus, 261, 263; *clavatus*, 261-263; *corrugis*, 263; *pistillaris*, 262, 263
Creolophus, 293; **agaricoides**, 294; **pulcherrimus**, 294; **septentrionalis**, 293
Creolophus, *Echinodontium*, *Gloiodon*, and *Hydnodon*, The genera. Type studies in the Hydnaceae—VI, 293
Crepidotus, 18, 26; *alveolus*, 27, 31, 32; **aquosus**, 30; **bicolor**, 28; *cacaophyllus*, 27; **calolepidoides**, 30; *calolepis*, 31; **cinchonensis**, 30; *Citri*, 27, 30; *croceosanguineus*, 28; *cuneiformis*, 29; *Dussii*, 28; **fumosifolius**, 31; *laceratus*, 29; *mollis*, 32; *musaeicola*, 27, 32; **parvulus**, 27; *Psychotriae*, 27; *pyrrhus*, 28, 29; **subcuneiformis**, 29; **substipitatus**, 31; **sulcatus**, 29, 30
Cryptodiscus araneo-cinctus, 248
Cryptoporus, 313
Cultures of heteroecious rusts, Further, 233
Cup-fungi, Some tropical, 185
Cylindrium, 45, 46, 55
Cyphellium, 118
Cystococcus, 129, 130, 145; *humicola*, 145, 146
Cytology of the Laboulbeniales, Faull's, 174
Dactylococcus, 125; *infusionum*, 118
Daedalea, 313; *quercina*, 115, 291
Dendrogaster, 313
Diaporthe Batatatis, 279; *parasitica*, 90
Diatrype patella, 249
Dicaeoma, 240-243
Dictyonema, 115, 118, 126
Dictyophora irpicina, 269
Didymascella, 7, 8; *Oxycedri*, 8, 9
Didymium anomalum, 250
Diplodia Akebiae, 248
Diploschistes, 129
Dodge, B. O., Faull's cytology of the Laboulbeniales, 174
Durand, E. J., The genus *Keithia*, 6
Eastern North America, The *Amanitas* of, 72
Echinodontium, 295, 296; *tinctorium*, 295
Echinodontium, *Gloiodon*, and *Hydnodon*, The genera *Creolophus*. Type studies in the Hydnaceae—VI, 293
Elfvigia megaloma, 291
Endocarpon pusillum, 122, 123
Endomyces scytonematum, 113
Endophyllum, 281, 282
Endothia, 274; *gyrosa*, 280; *parasitica*, 90, 274, 280; *radicalis*, 90, 280; *radicalis mississippiensis*, 280
Enerthenema syncarpon, 250
Entoloma, 259; *Grayanum*, 259; *lividum*, 259; **mirabile**, 68; *sinuatum*, 259
Ephebe, 112, 116, 124; *Hegetschweileri*, 113; *pubescens*, 109, 119
Epigloea bactrospora, 111, 118, 120
Erigeron, 242
Erysiphe graminis, 58
Eucalyptus, 35
Euglena viridis, 131

- Eusynchytrium*, 315
Evernia furfuracea, 130; *prunastri*, 149
Exobasidium, 88, 89
Exosporium Ulmi, 40
- Fagus*, 245
 Fairman, C. E., Notes on new species of fungi from various localities, 245
 Faull's cytology of the Laboulbeniales, 174
Favolus, 313
 Fink, Bruce, The nature and classification of lichens—II. The lichen and its algal host, 97
Flammula, 18; *areolata*, 36; *aureobrunnea*, 19; *aureoviridis*, 19; *bryophila*, 36; *chrysotrichoides*, 36; *depressa*, 36; *Earlei*, 36; *hispidella*, 36; *hypholomoides*, 36; *jalapensis*, 36; *lateritia*, 19; *Nashii*, 36; *olivacea*, 18; *palmicola*, 36; *parvula*, 36; *pholiotoides*, 36; *subpenetrans*, 36; *tenuis*, 36; *vinicolor*, 18
 Fleshy fungi, A bad year for, 315
Fomes, 295, 296, 313; *Laricis*, 287; *semitostus*, 39; *tinctorius*, 295; *ungulatus*, 291
 Fraser, W. P., Further cultures of heteroecious rusts, 233
Fraxinus, 247
Fuligo megaspora, 250
 Fungi, A bad year for fleshy, 315
 Fungi from various localities, Notes on new species of, 245
 Fungi, Illustrations of—XIII, 1; XIV, 93; XV, 257; XVI, 287
 Fungi, New species of, 67
 Fungi, Some tropical cup-, 185
 Fungus originate from pycnosporos or ascospores, A method of determining in analytic work whether colonies of the chestnut blight, 274
 Further cultures of heteroecious rusts, 233
Fusarium, 178, 180, 181, 279; *Solani*, 181; *vasinfectum*, 181
Fusarium problem, Wollenweber's studies on the, 178
Fusicladium Vanillae, 40
Fusidium, 45, 46
- Galera frustulenta*, 36
Ganoderma, 313
Gautieria, 313
Gaylussacia baccata, 238
 Genera *Creolophus*, *Echinodontium*, *Gloiodon*, and *Hydnodon*. Type studies in the Hydnaceae—VI, 293
 Genus *Hydnellum*, The. Type studies in the Hydnaceae—V, 194
 Genus *Keithia*, The, 6
 Genus *Phellodon*, The. Type studies in the Hydnaceae—IV, 62
 Genus *Pseudoplectania*, The, 299
 Genus *Sarcodon*, The. Type studies in the Hydnaceae—III, 12
 Genus *Synchytrium*, The, 315
Geopyxis aluticolor, 191; *elata*, 189, 190; *Mölderiana*, 192
Geotrichum, 45, 56; *candidum*, 56, 61; *cuboideum*, 56, 61
Gloeocapsa, 111, 124, 134; *polydermatica*, 118
Gloeocystis, 112
Gloeopeziza Rehmii, 112
Gloeosporium, 171; *Caryae*, 88
Gloiodon, 296, 298
Gloiodon, and *Hydnodon*, The genera *Creolophus*, *Echinodontium*. Type studies in the Hydnaceae—VI, 293
Glomerella, 171; *cingulata*, 171; *Gossypii*, 171; *lindemuthianum*, 171
Glomerella, Shear's studies of parasitic species of, 171
Glycophila, 45, 46, 57
Gnomonia Caryae, 88; *setacea macrospora*, 88
Gossypium, 59
Graphis, 105; *scripta*, 122, 123, 139
Grifola frondosa, 290
 Guide, Ridgway's new color, 172
Gymnoglossum, 313
Gymnophilus, 18, 21, 35, 36; *areolatus*, 24, 36; *aureobrunneus*, 19; *aureoviridis*, 19; *bryophilus*, 22, 36; *carbonarius*, 25; *chrysopellus*, 20, 23; *chrysotrichoides*, 21, 36; *chrysotrichus*, 21; *depressus*, 20, 36; *Earlei*, 22, 36; *helvoliceps*, 20; *hispidellus*, 24, 36; *hispidus*, 24; *hypholomoides*, 26, 36; *jalapensis*, 25, 36; *lateritius*, 19, 23; *Nashii*, 23, 36; *olivaceus*, 18; *palmicola*, 23, 36; *parvulus*, 19, 36; *penetrans*, 20, 26, 36; *pholiotoides*, 24, 36; *sapineus*, 26; *subpenetrans*, 20, 36; *tenuis*, 22, 36; *vinicolor*, 18
Gymnosporangium, 278
Gyroporus castaneus, 1
- Haematococcus pluvialis*, 131
Haematomma ventosum, 143
Halobysus, 45, 46
Haplochytrium, 315
 Harper, E. T., The identity of *Cantharellus brevipes* and *Cantharellus clavatus*, 261; The probable identity of *Stropharia epimyces* (Peck) Atk. with *Pilosace algeriensis* Fries, 167

- Heald, F. D., A method of determining in analytic work whether colonies of the chestnut blight fungus originate from pycnosporos or ascospores, 274
- Helicocephalum, 45, 46
- Helminthosporium, 279
- Helotium purpuratum, 192
- Hendersonia **coccolobina**, 247; **hy-pocarpa**, 246; Opuntiae, 38; Rosae, 247
- Heppia urceolata, 124
- Herpotrichia, 282; nigra, 282, 283
- Heteroecious rusts, Further cultures of, 233
- Heuchera cylindrica, 71
- Host, The lichen and its algal. The nature and classification of lichens—II, 97
- Hyalopsora, 237
- Hydnaceae, Type studies in the—III. The genus Sarcodon, 12; IV. The genus Phellodon, 62; V. The genus Hydnellum, 194; VI. The genera Creolophus, Echinodontium, Gloiodon, and Hydnodon, 293
- Hydnellum, 65, 198, 199, 204; **Dia-bolus**, 194; **geogenium**, 204; **hybridum**, 198; **inquinatum**, 202; **parvum**, 200; **Peckii**, 203; **Rickerii**, 201; sanguinarium, 196, 198; scrobiculatum, 196; suaveolens, 201, 202; velutinum, 196; **Ves-pertilio**, 198; zonatum, 199, 201
- Hydnellum, The genus. Type studies in the Hydnaceae—V, 194
- Hydnodon**, 297; **thelephorum**, 297
- Hydnodon, The genera Creolophus, Echinodontium, Gloiodon and. Type studies in the Hydnaceae—VI, 293
- Hydnofomes, 295, 296; tsugicola, 295
- Hydnophysa, 295, 296
- Hydnum, 288, 296, 298; acre, 13; agaricoides, 294; amicum, 62; boreale, 201; bubalinum, 14; carbunculus, 194, 195; cervinum, 14, 15; compactum, 201; conrescens, 199, 200; corrugatum, 293; cristatum, 13; cyathiforme, 64, 65, 199, 200; discolor, 294; ferrugineum, 195-198; fragile, 12; fragrans, 64; friabile, 294, 295; fuligineo-viola-ceum, 14; geogenium, 204, 205; graveolens, 62-64; hybridum, 197-199; imbricatum, 12, 14, 15; laevigatum, 14; leptopus, 62, 63; leptopus graveolens, 64; lateritium, 297, 298; melaleucum, 62-65; nigrum, 62, 198; Queletii, 198; parasiticum, 297; pulcherrimum, 294, 295; pullum, 62, 63; scrobiculatum, 196-198; septentrionale, 293; spongiosipes, 196; stratosum, 297; strigosum, 297; suaveolens, 201; sulphureum, 204, 205; thelephorum, 297; tomentosum, 64, 65; vellereum, 62; velutinum, 196, 197; Vespertilio, 198; zonatum, 62, 198-200
- Hymenochaete noxia, 39
- Hypheothrix Zenkeri, 112
- Hypholoma, 31; sublateritium, 26
- Hyphomycetes, Studies in North American—II, 45
- Hypocrea, 179
- Hypocreaceae, Maire's remarks on some, 176
- Hypodendrum, 18, 35; **scobifer**, 35
- Hypomyces, 176, 179, 180; armeniacus, 177; aureo-nitens, 177; boletinus, 176; chrysospermus, 176; hyalinus, 177; inaequalis, 177; macrosporus, 177; ochraceus, 177; papyraceus, 177; Solani, 181; tegillum, 177; violaceus, 177
- Hypophyllum album, 308
- Hysterangium, 313
- Hysterium, 105
- Identity of Cantharellus brevipes and Cantharellus clavatus, The, 261
- Identity of Stropharia epimyces (Peck) Atk. with Pilosace algeriensis Fries. The probable, 167
- Illustrations of fungi—XIII, 1; XIV, 93; XV, 257; XVI, 287
- Index to American mycological literature, 41, 91, 182, 251, 284, 317
- Inocybe, 224-226, 228, 229, 231; decipiens, 225; infelix, 224, 225; infida, 224-232; **minima**, 69
- Inocybe infida, Toxicological studies on the mushrooms Clitocybe illudens and, 224
- Inonotus hirsutus, 297
- Internal aecia, 303
- Irpex, 288
- Irpiciporus mollis, 288
- Ischnoderma, 313
- Jungermannia, 111; triophylla, 112
- Juniperus, 6-8, 11; communis, 8; Oxycedrus, 9
- Kalchbrennera, 268
- Karschia, 113, 114; destructans, 113
- Keithia, 6-8; tetraspora, 6-11; **thujina**, 6-9, 11; **Tsugae**, 7, 8, 10, 11
- Keithia, The genus, 6
- Laboulbenia chaetophora, 174; Gyrinidarum, 174
- Laboulbeniales, Faull's cytology of the, 174
- Lachnea crispata, 192; nigrella, 301

- Lactaria, 305; camphorata, 307; Chelidonium, 305; circellata, 307; deliciosa, 305; grisea, 306; insulsa, 306; mitissima, 307; mucida, 306; piperata, 307; scrobiculata, 305; subdulcis, 307; theiogala, 306; torminosa, 305; trivialis, 306; vellerea, 307, 308; zonaria, 306
- Lactariaeae of the Pacific Coast, The, 305
- Lactarius brevipes, 306; brevix, 306; deflexus, 306; exsuccus, 308; subserifluus, 307; villosus, 305; xanthogalactus, 307
- Laterna, 268
- Laudatea, 115, 126
- Leaia, 296
- Lecanora atrorufa, 127; caesiorufa, 127; coilocarpa, 127; ferruginea, 127; granatina, 118; pallida, 140; subfusca, 122, 125, 127
- Lecidea, 105; atrobrunnea, 140; enteroleuca, 122
- Lepiota, 94; cretacea, 85; procera, 257
- Leptogonium subtile, 119
- Leptonia **gracilipes**, 69, 70; **validipes**, 70
- Leptospora Musae, 89
- Lichen and its algal host, The. The nature and classification of lichens—II, 97
- Lichens, The nature and classification of—II. The lichen and its algal host, 97
- Lichina, 138; confinis, 124; pygmaea, 119
- Liquidambar, 295
- Liriodendron, 54
- Literature, Index to American mycological, 41, 91, 182, 251, 284, 317
- Lycoperdon giganteum, 52
- Lycopus virginicus, 303, 304
- Lysurus, 268
- Maire's remarks on some Hypocreaceae, 176
- Malbranchea, 45, 57; pulchella, 57; **pulveracea**, 57, 61
- Mangifera indica, 247
- Marasmius, 249
- Melampsora arctica, 238, 239; Medusae, 238, 239
- Melanoleuca, 206; **anomala**, 214, 223; **arenicola**, 214, 223; **avellanea**, 215, 223; **avellaneifolia**, 215, 223; **bicolor**, 215, 223; **californica**, 216, 223; **collybiiformis**, 216, 223; **dryophila**, 217, 220, 223; **farinacea**, 217, 223; **Harperi**, 217, 223; **nuciolens**, 218, 223; **Olesonii**, 218, 223; **oreades**, 218, 223; **pini-cola**, 219, 223; **platyphylla**, 219, 223; **portolensis**, 219, 223; **rosei-brunnea**, 216, 220, 223; **rudericola**, 220, 223; **secedifolia**, 221, 223; **striatella**, 221, 223; **sublurida**, 221, 223; **submulticeps**, 221, 223; **subpessundata**, 217, 222, 223; **subvelata**, 222, 223; **tenuipes**, 223
- Melanoleuca, New species of Clitocybe and. The Agaricaceae of the Pacific Coast—IV, 206
- Melascypha, 299; melaena, 300
- Merulius clavatus, 263; lacrymans, 89
- Method of determining in analytic work whether colonies of the chestnut blight fungus originate from pycnosporos or ascospores, A, 274
- Michelia fuscata, 120
- Microsphaera Platani, 58
- Monilia, 45-47; Aspergillus, 47; aurantiaca, 48; aurea, 47; aureofulva, 48; Avenae, 59; candida, 47, 52, 60; capitata, 47; Cerasi, 51; cespitosa, 47; cespitosa aurea, 47; cinerea, 50; crustacea, 47; diffusa, 60; divaricata, 47; effusa, 48; fructigena, 50; fungicola, 51; fusconigra, 60; glauca, 47; globosa, 60; Harknessii, 59; hyalina, 57, 58; Linhartiana, 51, 60; Martinii, 52; megalospora, 49; mycophila, 60; nidulans, 47; Peckiana, 52, 60; pencillata, 60; pulveracea, 57; punctans, 60; racemosa, 47; ramosa, 47; rosea, 47; rubiginosa, 60; simplex, 47; sitophila, 52; urediniformis, 60; viridi-flava, 60
- Mucor Aspergillus, 47; crustaceus, 47
- Murrill, W. A., A bad year for fleshy fungi, 316; Illustrations of fungi—XIII, 1; XIV, 93; XV, 257; XVI, 287; Sterility in Pholiota candicans (Bull.) Schroet., 314; The Agaricaceae of the Pacific Coast—IV. New species of Clitocybe and Melanoleuca, 206; The Agaricaceae of tropical North America—IV, 18; The Amanitas of eastern North America, 72
- Mushrooms Clitocybe illudens and Inocybe infida, Toxicological studies on the, 224
- Mycena, 36
- Mycobacidia, 105
- Mycobilimbia, 105
- Mycocalicium, 105
- Mycological literature, Index to American, 41, 91, 182, 251, 284, 317
- Mycosphaerella lageniformis, 249
- Naevia canadica, 249

- Nature and classification of lichens, The—II. The lichen and its algal host, 97
- Nectandra, 22
- Nectria, 112, 179, 180; Ipomoeae, 180; phycophila, 112; Solani, 181; Vannillae, 40
- Nectriopsis, 176, 177, 179
- Neocosmospora, 181; vasinfecta, 181
- Neopeckia, 282; Coulteri, 282, 283
- New color guide, Ridgway's, 172
- New species of Clitocybe and Melanoleuca. The Agaricaceae of the Pacific Coast—IV, 206
- New species of fungi, 67
- New species of fungi from various localities, Notes on, 245
- News and notes, 37, 87, 249, 278, 312; and reviews, 170
- Nigredo, 240–242
- North America, The Agaricaceae of tropical—IV, 18
- North America, The Amanitas of eastern, 72
- North America, Uredinales on Carex in, 240
- North American Hyphomycetes, Studies in—II, 45
- Nostoc, 115, 119, 120, 124, 125, 130, 134, 136–138; lichenoides, 118
- Notes, News and, 37, 87, 249, 278, 312; and reviews, 170
- Notes on new species of fungi from various localities, 245
- Ocellaria Vanillae, 40
- Oidium, 45, 47, 48; albipes, 59; Asteris-punicel, 59; aureum, 48, 61; candidum, 59; compactum, 58; corticale, 60; erysiphoides, 59; Euonymi-japonici, 58; fructigenum, 50; inquinans, 60; irregulare, 60; Lactis, 52, 53; leucoconium, 58; leucogonium, 58; leuconium, 58; megalosporum, 48, 49, 61; **Murrilliae**, 48, 49, 61; obductum, 58; pirinum, 58; simile, 48, 61; Tuckeri, 58
- Ombrophila limosa, 249
- Omphalaria, 124
- Onoclea, 237; sensibilis, 234, 236, 237, 239; Struthiopteris, 234–236, 239
- Oospora, 45, 49, 53, 61; **Arthurii**, 50, 52, 61; candida, 50, 52; candidula, 55; **Cerasi**, 50, 51, 61; **cinerea**, 50, 61; crustacea, 45; cuboidea, 56; cucumeris, 60; fructigena, 50, 61; **fungicola**, 50, 51, 61; heterospora, 60; hyalinula, 55; Lactis, 45, 53; **Linhartiana**, 50–52, 61; **Martinii**, 50, 52, 61; Nicotianae, 54; pallida, 60; scabies, 61; similis, 48; Tulipiferae, 54
- Oosporoidea**, 52; **Lactis**, 53
- Opegrapha, 120, 124; filicina, 124; subsiderella, 122, 123; varia, 119, 120, 124; vulgata, 127
- Ophionectria, 179
- Oplismenus hirtellus, 246
- Oscillatoria, 125
- Osmunda, 235; Claytoniana, 234, 235, 239
- Otidea domingensis, 192
- Otidella, 299; fulgens, 302; **nigrella**, 301
- Ovularia, 61; isarioides, 61
- Pacific Coast, The Agaricaceae of the—IV. New species of Clitocybe and Melanoleuca, 206
- Pacific Coast, The Lactariae of the, 305
- Paepalopsis, 45, 46
- Palmella, 111, 112; botryoides, 111, 117, 118, 120
- Panaeolus epimyces, 167, 168
- Pannaria brunnea, 138; muscorum, 124; nigra, 120; tryptophylla, 118, 124
- Parasitic species of Glomerella, Shear's studies of, 171
- Parmelia acetabulum, 127, 129; aspidota, 133; olivacea, 152; physodes, 148
- Parüphrädria, 111; Heimerlii, 111
- Patinella, 105
- Peck, C. H., New species of fungi, 67
- Peckiella, 176
- Peltandra virginica, 304
- Peltigera, 131; canina, 130
- Penicillium, 47
- Peridermium, 235; balsameum, 233–237; Peckii, 11, 238
- Perisporium Wrightii, 38
- Pertusaria, 129; communis, 122, 125, 129
- Pestalotia, 245; truncata Rubi, 245; truncata **septoriana**, 245
- Pezicula eximia, 249
- Peziza, 187; Afzelii, 189; aluticolor, 191; Colensoi, 191; crispata, 192; cyanoderma, 302; domingensis, 186, 192; fulgens, 302; fuscocana, 300, 301; Harmoge, 192; Hindsii, 186, 188, 189; Hystrich, 186, 188; insititia, 190; melaena, 300; melania, 300, 302; **nigrella**, 299, 301; onotica, 186; spongiosa, 300, 302; striispora, 186, 188, 189; stygia, 301; **sulcipes**, 188–190; Tricholoma, 187–189; vogesiaca, 300, 302
- Phacidium tetrasporum, 6, 8
- Phaeolus, 313
- Phallogaster, 313
- Phegopteris Dryopteris, 234–236, 239

- Phellinus, 313
 Phellodon, 65, 66, 298; **amicus**, 62-64; **carnosus**, 65; **niger**, 62; **pul-lus**, 62-64; **tomentosus**, 64
 Phellodon, The genus. Type studies in the Hydnaceae—IV, 62
 Phillipsia, 186, 192; **domingensis**, 192, 193; **kermesina**, 192; **subpurpurea**, 192
 Pholiota, 18, 21, 25, 32, 35, 39; **avellanea**, 32; **Brittoniae**, 35; **Broadwayi**, 32; **bryophila**, 33; **cinchonensis**, 33; **cubensis**, 34; **martinicensis**, 34; **Musae**, 34; **praecox**, 314; **unicolor**, 33; **ventri-cosa**, 35
 Pholiota candicans (Bull.) Schroet., Sterility in, 314
 Pholiotina Musae, 34
 Phoma Batatae, 279; **pigmentivora**, 40; **rhodocarpa**, 246
 Phyllactidium, 124, 125, 136
 Phyllosticta **Mortoni**, 247; **Vanillae**, 40
 Physalis, 264
 Physcia, 130; **apiolia**, 137; **ciliaris**, 125; **pulverulenta**, 125, 131; **stel-laris**, 127, 148, 152
 Physma, 137; **chalazanum**, 124
 Pilocratera, 185, 187; **Engleriana**, 189, 190; **Hindsii**, 189; **Tricholoma**, 188
 Pilosace algeriensis, 167-169
 Pilosace algeriensis Fries, The prob-able identity of Stropharia epimyces (Peck) Atk. with, 167
 Pinus Strobilus, 67
 Piptoporus, 313
 Pirus coronaria, 58
 Placodium murorum, 129, 132
 Placynthium, 119
 Platanus orientalis, 59, 61
 Plectania nigrella, 301
 Pleurococcus, 125, 127, 129, 130, 145; **vulgaris**, 118, 130, 131
 Pleurotus ulmarius, 316
 Poa pratensis, 264
 Podostroma, 179
 Polyblastia rugulosa, 122, 123
 Polychidium muscicolum, 138
 Polycoccus punctiformis, 118, 138
 Polyporus, 88, 313, 314; **albo-sordes-cens**, 88; **lucidus**, 115; **rufopodex**, 88
 Polyscytalum, 45, 46, 55; **cylindroides**, 55; **fecundissimum**, 55; **sericeum**, 55, 56
 Polystictus, 313
 Populus fastigiata, 245; **grandiden-tata**, 238, 239
 Porodisculus, 313
 Poronidulus conchifer, 289
 Probable identity of Stropharia epi-myces (Peck) Atk. with Pilosace algeriensis Fries, The, 167
 Propolidium, 7; **Tsugae**, 10
 Propolis faginea, 7
 Protococcus, 127
 Protoglossum, 313
 Protuberia, 313
 Prunus, 51
 Psathyra, 36
 Pseudoplectania, 299; **fulgens**, 299, 302; **melaena**, 300; **melania**, 299; **nigrella**, 299, 301, 302; **stygia**, 299, 301; **vogesiacae**, 299, 300, 302
 Pseudoplectania, The genus, 299
 Psidium, 19
 Psilocybe uda, 314
 Psora ostreata, 150
 Psychotria glabrata, 27
 Ptychogaster, 314
 Puccinia, 9; **angustata**, 303, 304; **car-i-cina**, 240; **Caricis**, 240; **graminis**, 240, 304; **Peckiana**, 281; **striato-spora**, 71
 Pucciniastrum Myrtilli, 237-239
 Pulparia spongiosa, 300
 Pycnochytrium, 315
 Pycnosporos or ascospores, A method of determining in analytic work whether colonies of the chestnut blight fungus originate from, 274
 Pyrenochaeta **fraxinina**, 247
 Pyrenula nitida, 120, 124
 Pyxidiophora, 176
 Quercus, 58; **alba**, 58
 Racoblenna, 119
 Ramalina, 133; **calicaris**, 125
 Ramularia, 180
 Raphidium polymorphum, 131
 Remarks on some Hypocreaceae, Maire's, 176
 Reviews, News, notes and, 170
 Rhinotrichum, 48; **Curtisii**, 48; **pul-veraceum**, 57
 Rhizocarpon, 114; **alboatrum**, 114; **geographicum**, 114
 Rhymbocarpus, 114; **punctiformis**, 114
 Ribes, 242
 Ricker, P. L., Ridgway's new color guide, 172
 Ridgway's new color guide, 172
 Rinodina sophodes, 127, 148
 Rivularia, 119, 138; **nitida**, 118, 119
 Roccella, 124, 140; **phycopsis**, 124
 Rosa villosa, 247
 Rostkovites granulatus, 3; **subaureus**, 3
 Rubus, 245
 Russula, 305, 308; **abietina**, 309; **albi-dula**, 310; **alutacea**, 310; **azurea**, 311; **bicolor**, 311; **brevipes**, 308; **chameleontina**, 309; **crenulata**,

- 310; *delica*, 308; *deliciosa*, 308; *drimeja*, 309; *emetica*, 76, 309; *flaviceps*, 310; *granulata*, 309; **Murrillii**, 310; *nigrescens*, 308; *nigricans*, 308; *obscura*, 309; *pectinata*, 311; Turci, 309; *veternosa*, 309
 Rusts, Further cultures of heteroecious, 233
Salix, 238, 239
Sarcodon, 12; *acre*, 13; *fuligineo-violaceus*, 13; **fumosus**, 16; *imbriatus*, 14; *laevigatus*, 14; **Murrillii**, 15; *radicatus*, 13; *reticulatus*, 12; **roseolus**, 16
Sarcodon, The genus. Type studies in the Hydnaceae—III, 12
Sarcoscypha, 185, 186; *coccinea*, 186; *Colensoi*, 191; *striispora*, 188
Scenedesmus obtusus, 131
Scirpus, 303
Sclerodon, 296
Sclerotina, 109; *fructigena*, 50; *tuberosa*, 109
Scutiger, 290; *griseus*, 289
Scytonema, 113, 118, 124–126, 136, 138; *myochrous*, 109
 Seaver, F. J., Maire's remarks on some Hypocreaceae, 176; Shear's studies of parasitic species of *Glomerella*, 171; Some tropical cup-fungi, 185; The genus *Pseudoplectania*, 299; Wollenweber's studies on the *Fusarium* problem, 178
Septoria, 245; **Carricerae**, 246
Seurattia Coffeicola, 40; *Vanillae*, 40
 Shear's studies of parasitic species of *Glomerella*, 171
Simblum, 267–269, 271, 273; *australe*, 265; *Lorentzii*, 265; *periphragmoides*, 266, 269; *pilidiatum*, 265; *rubescens*, 265; *sphaerocephalum*, 264, 266, 267, 269–272
Simblum sphaerocephalum, The structure of, 264
Sirosiphon, 116, 119, 124, 136, 138; *alpinus*, 109; *pulvinatus*, 109, 118
 Smith, C. S., Clark, E. D., & Toxicological studies on the mushrooms *Clitocybe illudens* and *Inocybe infida*, 224
Solidago, 242
Solorina crocea, 118
 Some tropical cup-fungi, 185
 Species of *Clitocybe* and *Melanoleuca*, New. The Agaricaceae of the Pacific Coast—IV, 206
 Species of fungi from various localities, Notes on new, 245
 Species of fungi, New, 67
 Species of *Glomerella*, Shear's studies of parasitic, 171
Sphaerella Opuntiae, 38
Sphaeria Lactifluorum, 179
Sphaeropsis Cocolobae, 246; **rhodocarpa**, 246
Sphaerotheca pannosa, 58
Sphagnum, 300–302
Spilonema paradoxum, 119
Sporendonema, 45; *Casei*, 45; *myophilum*, 46; *terrestre*, 45
Steccherinum, 293
Stereocaulon, 118
 Sterility in *Pholiota candicans* (Bull.) Schroet., 314
Stichococcus bacillaris, 115, 131, 132; *bacillaris fungicola*, 115
Stictis, 7; *Tsugae*, 7, 10, 11
Stigeoclonium tenue, 131
Stigmatomma cataleptum, 121
Stigonema mammosum, 109
Strigula, 120, 151; *complanata*, 120
Stropharia, 167, 168; *coprinophila*, 167, 168; *epimyces*, 168
Stropharia epimyces (Peck) Atk. with *Pilosace algeriensis* Fries, The probable identity of, 167
 Structure of *Simblum sphaerocephalum*, The, 264
 Studies in North American Hyphomycetes—II, 45
 Studies in the Hydnaceae, Type—III. The genus *Sarcodon*, 12; IV. The genus *Phellodon*, 62; V. The genus *Hydnellum*, 194; VI. The genera *Creolophus*, *Echinodontium*, *Gloiodon*, and *Hydnodon*, 293
 Studies of parasitic species of *Glomerella*, Shear's, 171
 Studies on the *Fusarium* problem, Wollenweber's, 178
 Studies on the mushrooms *Clitocybe illudens* and *Inocybe infida*. Toxicological, 224
 Sumstine, D. R., Studies in North American Hyphomycetes—II, 45
Synalissa, 124
Synchytrium, 313; *Amsinckiae*, 313; *andinum*, 313; *aureum*, 315; *innominatum*, 313; *papillatum*, 313; *pluriannulatum*, 315; *Succisiae*, 315; *Taraxaci*, 315
Synchytrium, The genus, 315
Taphrina, 88; *lapponica*, 89
Thelephora, 105; *padinaeformis*, 297, 298
Thelotrema lepadinum, 122
Thuja, 6–8, 11; *occidentalis*, 10
Thyridaria tarda, 39
Tilia, 294

- Torula*, 45, 53, 60; *aurea*, 48; *fructigena*, 49, 50; *pallida*, 60
Toruloidea, 53, 60; ***candidula***, 53, 55, 61; ***effusa***, 53, 61; **Nicotianae**, 53, 54; ***Tulipiferae***, 53, 54, 61; ***Unangstii***, 53, 54, 61
 Toxicological studies on the mushrooms *Clitocybe illudens* and *Inocybe infida*, 224
Trametes, 313; *Pini*, 115
Trentepohlia, 110, 116, 120, 124, 125, 127, 132, 136, 138-140, 147; *aurea*, 115, 116; *umbrina*, 109, 118, 119
Trichoderma aureum, 47
Tricholoma, 206, 208; *album*, 207, 220; *anomalum*, 223; *arenicola*, 223; *avellaneifolium*, 223; *avellaneum*, 223; *bicolor*, 223; *californicum*, 223; *collybiiforme*, 223; *dryophilum*, 223; *farinaceum*, 223; *Harperi*, 223; *luridum*, 221; *maculatescens*, 209; *nuciolens*, 223; *nudum*, 208; *Olesonii*, 223; *oreades*, 223; *pes-sundatum*, 222; *pinicola*, 223; *platyphyllum*, 223; *portolense*, 223; *roseibrunneum*, 223; *ru-dericola*, 223; *secedifolium*, 223; *striatellum*, 223; *subluridum*, 223; *submulticeps*, 223; *sub-pessundatum*, 223; *subvelatum*, 223; *tenuipes*, 223
Trichoscypha, 185, 187; *Afzelii*, 189; *Hindsii*, 189; *insititia*, 190; *sul-cipes*, 189; *Tricholoma*, 188
 Tropical cup-fungi, Some, 185
 Tropical North America, The *Agari-caceae* of—IV, 18
Tsuga, 6-8, 11, 233; *canadensis*, 10, 237-239
 Type studies in the *Hydnaceae*—III. The genus *Sarcodon*, 12; IV. The genus *Phellodon*, 62; V. The genus *Hydnellum*, 194; VI. The genera *Creolophus*, *Echinodontium*, *Gloio-don*, and *Hydnodon*, 293
Ulothrix, 125
Uncinula necator, 58
 Uredinales on *Carex* in North Amer-ica, 240
Uredinopsis, 223, 237, 304; *Atkin-sonii*, 236, 239; *mirabilis*, 236, 239; *Osmundae*, 235, 239; *Phegopteridis*, 236, 239; *Struthiopteridis*, 234, 235, 239
Uredo Scabies, 40
Uromyces Caladii, 304
Urophlyctis pluriannulatus, 313
Urtica, 242
Vaccinium canadense, 237-239; *penn-sylvanicum*, 52, 238
Vaginata, 81, 94; *agglutinata*, 82-85, 94; *albocreata*, 79, 82, 84; *fari-nosa*, 82-85; *livida*, 82; *parcivol-vata*, 82, 83; *plumbea*, 82, 83, 85, 86; *plumbea alba*, 84; *plumbea strangulata*, 82; *pusilla*, 82, 83; *vaginata*, 82
Valsaria Akebiae, 248
Vaucheria sessilis, 127
Venenarius, 72, 81, 94, 258; **Caesar-eus**, 72, 73, 85; ***crenulatus***, 73, 77; ***cothurnatus***, 72, 74, 77, 79, 81, 96; ***flavorubescens***, 73, 76, 95; ***Frostianus***, 73, 76, 79, 96; ***junquilleus***, 80; ***Morrisii***, 73, 75; ***muscarius***, 73, 75-77, 94, 96; ***pantherinus***, 80, 81, 84; ***phal-loides***, 72, 74, 79-81, 93; ***por-phyrius***, 81; ***recutitus***, 81; ***rubens***, 73, 75, 76, 79, 95; ***rus-suloides***, 73, 76, 77, 80; ***solitarius***, 73, 77, 79, 85; ***spissus***, 81; ***spretus***, 72, 73, 81; ***velatipes***, 73, 75
Verrucaria, 124; *muralis*, 123, 127
Verticillium, 180
 Wilson, G. W., The genus *Synchy-trium*, 315
 Wolf, F. A., Internal aecia, 303
 Wollenweber's studies on the *Fu-sarium* Problem, 178
 Wynnea, 186
Xanthoria, 128; *parietina*, 125, 127-132, 151
Xylaria, 178
Zea Mays, 52
Zygodesmus, 60